

Spring 5-10-2016

A carbon dioxide stripping model for mammalian cell culture in manufacturing scale bioreactors

Zizhuo Xing

Bristol-Myers Squibb Company, zizhuo.xing@bms.com

Nicholas Abuabsi

Bristol-Myers Squibb Company

Michael Borys

Bristol-Myers Squibb Company

Zheng Jian

Bristol-Myers Squibb Company

Follow this and additional works at: http://dc.engconfintl.org/cellculture_xv



Part of the [Biomedical Engineering and Bioengineering Commons](#)

Recommended Citation

Zizhuo Xing, Nicholas Abuabsi, Michael Borys, and Zheng Jian, "A carbon dioxide stripping model for mammalian cell culture in manufacturing scale bioreactors" in "Cell Culture Engineering XV", Robert Kiss, Genentech Sarah Harcum, Clemson University Jeff Chalmers, Ohio State University Eds, ECI Symposium Series, (2016). http://dc.engconfintl.org/cellculture_xv/26

This Abstract is brought to you for free and open access by the Proceedings at ECI Digital Archives. It has been accepted for inclusion in Cell Culture Engineering XV by an authorized administrator of ECI Digital Archives. For more information, please contact franco@bepress.com.

A CARBON DIOXIDE STRIPPING MODEL FOR MAMMALIAN CELL CULTURE IN MANUFACTURING SCALE BIOREACTORS

Zizhuo Xing, Bristol-Myers Squibb Company
zizhuo.xing@bms.com

Nicholas A. Abuabsi, Bristol-Myers Squibb Company

Michael C. Borys, Bristol-Myers Squibb Company

Zheng Jian Li, Bristol-Myers Squibb Company

Achieving adequate CO₂ stripping rates in large scale bioreactors is an important consideration during the scale up of animal cell cultures to large scale bioreactors due to the use of relatively low power input and gas sparging rates. It has previously been reported that cell growth, productivity, and product quality attributes such as glycosylation can be significantly impacted when cells are exposed to high CO₂ environments. CO₂ stripping models that depend on the CO₂ mass transfer coefficient have been applied to simulate CO₂ profiles in cell cultures using varied sparger types, reagents for pH adjustment, gas flow rates, and agitation speeds. These models were reported as being validated for a cell culture after cell exponential growth phase. However, in recent years, cell culture processes have been improved to enhance productivity in part through a longer exponential growth phase to achieve higher viable cell densities, making those models less relevant. The current CO₂ stripping models were tested in several improved cell culture processes and resulted in predicted CO₂ profiles not fitting the measured CO₂ profiles. A modified CO₂ stripping model was then developed, of which CO₂ stripping is independent of the CO₂ mass transfer coefficient. Instead, CO₂ stripping is a function of gas flow rates, the residence time of bubbles in the liquid, the time of bubbles being saturated with CO₂, and CO₂ concentrations. The model was validated with two CHO cell culture processes that achieved different peak viable cell density (approximately 7×10^6 cells/mL and 12×10^6 cells/mL) in 25,000-L and 5,000-L manufacturing bioreactors, respectively. The CO₂ stripping model was also applied to optimize cell culture conditions to reduce CO₂ level in cell cultures in the manufacturing scale bioreactors.