

Spring 5-10-2016

# Improving the metabolic efficiency of mammalian cells and its impact on glycoproteins quality

Eric Karengera

*Ecole Polytechnique de Montreal, eric.karengera@polymtl.ca*

Gregory De Crescenzo

*Ecole Polytechnique de Montreal*

Yves Durocher

*National Research Council Canada*

Olivier Henry

*Ecole Polytechnique de Montreal, olivier.henry@polymtl.ca*

Follow this and additional works at: [http://dc.engconfintl.org/cellculture\\_xv](http://dc.engconfintl.org/cellculture_xv)



Part of the [Biomedical Engineering and Bioengineering Commons](#)

---

## Recommended Citation

Eric Karengera, Gregory De Crescenzo, Yves Durocher, and Olivier Henry, "Improving the metabolic efficiency of mammalian cells and its impact on glycoproteins quality" in "Cell Culture Engineering XV", Robert Kiss, Genentech Sarah Harcum, Clemson University Jeff Chalmers, Ohio State University Eds, ECI Symposium Series, (2016). [http://dc.engconfintl.org/cellculture\\_xv/101](http://dc.engconfintl.org/cellculture_xv/101)

This Abstract is brought to you for free and open access by the Proceedings at ECI Digital Archives. It has been accepted for inclusion in Cell Culture Engineering XV by an authorized administrator of ECI Digital Archives. For more information, please contact [franco@bepress.com](mailto:franco@bepress.com).

## IMPROVING THE METABOLIC EFFICIENCY OF MAMMALIAN CELLS AND ITS IMPACT ON GLYCOPROTEINS QUALITY

Eric Karengera, École Polytechnique de Montréal, Canada  
eric.karengera@polymtl.ca

Gregory De Crescenzo, École Polytechnique de Montréal, Canada  
Yves Durocher, Human Health Therapeutics Portfolio, National Research Council, Canada

Olivier Henry, École Polytechnique de Montréal, Canada  
P.O. Box 6079, Centre-ville station, Montréal, Québec H3C 3A7, Canada  
T: 514-340-4711 (x2191) F: 514-340-4159, olivier.henry@polymtl.ca

Key Words: HEK293, Fed-batch, protein glycosylation, PYC-overexpression.

Glycosylation is a critical quality attribute for recombinant therapeutic proteins, which can be impacted by a number of process conditions, including waste metabolite accumulation. While fed-batch strategies that consist in substituting or controlling the main substrates at low concentrations have proven generally effective at improving protein titer, they can also adversely affect product glycosylation. Metabolic engineering strategies aiming at reducing by-product formation may thus be beneficial for ensuring product quality consistency. In this work, we have specifically investigated the impact of PYC2 overexpression on the quality of a recombinant glycoprotein of therapeutic interest, the interferon  $\alpha 2b$  (IFN $\alpha 2b$ ) that has one O-glycosylation site. To this end, batch and fed-batch cultures were performed and product characteristics were measured for both the PYC expressing HEK293 clone and the parental cells.

SDS-PAGE and Western Blot analysis of batch culture harvests revealed two distinct bands corresponding to glycosylated and non-glycosylated fractions of IFN $\alpha 2b$ , as subsequently confirmed via SDS-PAGE analysis of purified samples loaded along with a non-glycosylated commercial standard produced in *E.coli*. As inferred from densitometry analysis of the gels, the cultures with PYC-expressing cells were shown to sustain a significantly higher percentage of glycosylated IFN $\alpha 2b$  at the late stage of the culture, which was correlated with the prolonged viability and reduced accumulation of waste metabolites. Differences between the two cell lines in terms of cell viability and protein quality were even more pronounced when performing fed-batch cultures during which glucose was maintained at high levels. To investigate the potential impact of ammonia, batch cultures with various glutamine substitutes were also performed. Among the different substitutes tested, pyruvate led to the lowest ammonia production with no significant impact on protein titer. Of salient interest, the results suggest that substituting glutamine by  $\alpha$ -ketoglutarate, glutamate or pyruvate may allow to maintain a higher fraction of glycosylated proteins during late-stage batch cultures.