

Spring 5-13-2016

Improvement of CHO specific productivity using amino acid derivatives

Aline Zimmer

Merck, aline.zimmer@merckgroup.com

Caroline Hecklau

Merck

Sascha Pering

Merck

Alisa Schnellbaecher

Merck

Thomas Eichhorn

Merck

Follow this and additional works at: http://dc.engconfintl.org/cellculture_xv

 Part of the [Biomedical Engineering and Bioengineering Commons](#)

Recommended Citation

1. Zeck, A., et al., Low level sequence variant analysis of recombinant proteins: an optimized approach. PLoS One, 2012. 7(7): p. e40328. 2. Khetan, A., et al., Control of misincorporation of serine for asparagine during antibody production using CHO cells. Biotechnol Bioeng, 2010. 107(1): p. 116-23. 3. Feeney, L., et al., Eliminating tyrosine sequence variants in CHO cell lines producing recombinant monoclonal antibodies. Biotechnol Bioeng, 2013. 110(4): p. 1087-97. 4. Kshirsagar, R., et al., Controlling trisulfide modification in recombinant monoclonal antibody produced in fed-batch cell culture. Biotechnol Bioeng, 2012. 109(10): p. 2523-32. 5. Zimmer, A., et al., Improvement and simplification of fed-batch bioprocesses with a highly soluble phosphotyrosine sodium salt. J Biotechnol, 2014. 186: p. 110-118.

IMPROVEMENT OF CHO SPECIFIC PRODUCTIVITY USING AMINO ACID DERIVATIVES

Aline Zimmer, Merck KGaA, Germany
Aline.Zimmer@merckgroup.com

Caroline Hecklau, Merck KGaA, Germany
Sascha Pering, Merck KGaA, Germany
Alisa Schnellbaecher, Merck KGaA, Germany
Thomas Eichhorn, Merck KGaA, Germany

Key words: CHO productivity, amino acid derivatives, fed-batch, IgG quality attributes

Industrial fed-batch cultivation of mammalian cells is used for the production of therapeutic proteins such as monoclonal antibodies. Besides medium ensuring initial growth, feeding is necessary to improve growth, viability and antibody production. Amino acids are key elements for the cellular metabolism but also for the quality of the recombinant protein produced. Several studies have already demonstrated the link between amino acid availability and sequence variants [1-3] or specific modifications like trisulfide bonds [4]. To avoid such modifications, the chemical modification of amino acids is an interesting alternative to modulate their overall solubility [5], stability or chemical reactivity.

In this study, we analyzed the beneficial effect of using amino acid derivatives in neutral pH feeds on CHO cell growth and specific productivity in small scale and bioreactor experiments. The mechanisms of extra- or intracellular amino acid metabolization were investigated as well as the interaction with other media or cellular components. Additionally, gene expression arrays and western blot analyses were used to decipher the mechanisms of increased productivity. Finally, LC-MS based methods were used to study the impact of the chemical modification on several IgG critical quality attributes.

1. Zeck, A., et al., *Low level sequence variant analysis of recombinant proteins: an optimized approach*. PLoS One, 2012. **7**(7): p. e40328.
2. Khetan, A., et al., *Control of misincorporation of serine for asparagine during antibody production using CHO cells*. Biotechnol Bioeng, 2010. **107**(1): p. 116-23.
3. Feeney, L., et al., *Eliminating tyrosine sequence variants in CHO cell lines producing recombinant monoclonal antibodies*. Biotechnol Bioeng, 2013. **110**(4): p. 1087-97.
4. Kshirsagar, R., et al., *Controlling trisulfide modification in recombinant monoclonal antibody produced in fed-batch cell culture*. Biotechnol Bioeng, 2012. **109**(10): p. 2523-32.
5. Zimmer, A., et al., *Improvement and simplification of fed-batch bioprocesses with a highly soluble phosphotyrosine sodium salt*. J Biotechnol, 2014. **186**: p. 110-118.